


Observing Asteroidal Occultations from Multiple Stations

2012 August 25

ESOP-32, Barcelona, Cataluña, Spain

David W. Dunham, IOTA

Geometry of an Asteroid Occultation




Remote Stations for Asteroidal Occultations

- Separation should be many km, much larger than for grazes, so tracking times & errors are too large
- Unguided is possible since the prediction times are accurate enough, to less than 1 min. = $\frac{1}{4}^\circ$
- Point telescope beforehand to same altitude and azimuth that the target star will have at event time and keep it fixed in that direction
- Plot line of target star's declination on a detailed star atlas; I used the Millennium Atlas, but now use Guide 8 to produce the charts
- From the RA difference and event time for the area of observation, calculate times along the declination line
- Adjust the above for sidereal rate that is faster than solar rate, add 10 seconds for each hour before the event
- Can usually find “guide stars” that are easier to find than the target
- Find a safe but accessible place for both the attended & remote scopes
- Separation distance limited by travel time & tape to start tapes, but we have had some success with programmable remote control devices to turn on the recordings; then the only limit is battery life, which can be several hours
- Roger Venable uses VCR's with timed starts, allows larger separation
- Sometimes it is better to have remote sites attended for starting equipment later (allows larger separations) and security

Occultation of the 6.0-mag. Close Double Star SAO 78349 by (9) Metis on 2001 September 7

- The star was known to be a close double, sep. about $0.08''$ with 6.5 and 6.9-mag. Components, from a photoelectric lunar occultation recording at McDonald Obs., Texas, on 1973 April 9
- Best asteroidal occultation of 2001 in the U.S.A.
- Unfortunately, 1 night before the occultation of a 7th-mag. Star by Uranus' satellite Titania in Europe & n. S. America
- I made the first REMOTE recording of an asteroidal occultation during this event, in the Sacramento Valley of northern California
- Kent Okasaki tried a remote observation of this event, but he tried to track with a 20cm SCT, and the tracking wasn't accurate enough

Sky-plane plot of Metis occ'n from March 2002 S&T



Remote equipment at Orland, CA



Another view


This used my image intensifier and a 50mm Nikon lens, but similar results (with a narrower, about 3° , field of view) are possible with the PC164C.



Successful Remote + Attended Positive Observations from 2 or more stations, 2001 to 2008

- 2001 Sept. 7, 9 Metis, northern California, D. Dunham
- 2002 April 21, Oriola, Washington, S. Preston
- 2003 Jan. 17, Bathilde, Georgia, R. Venable
- 2004 July 1, Nanon, s. Calif., D. Dunham, but D. Stockbauer was at “remote” site, turned on recorder without changing pointing
- 2004 Oct. 6, Ute, North Carolina, D. Dunham
- 2004 Oct. 29, Flora, New Mexico, D. Dunham
- 2005 Mar. 12, Bathseba, Georgia, R. Venable
- 2005 May 13, Dufour, New South Wales (AU), D. Gault (home “remote” & mobile)
- 2005 Dec. 1, Laurentia, Georgia, R. Venable (**first time, two successful multiple deployments in one night**)
- 2005 Dec. 1, Dike, Maryland & Virginia, D. Dunham (**3 positives, star close double**)
- 2005 Dec. 3, Europa, California, D. Dunham
- 2006 Jan. 28, Veritas, North Carolina, D. Dunham
- 2006 Feb. 24, Turandot, Indiana, D. Dunham
- 2006 Feb. 26, Abnoba, Florida, R. Venable
- 2006 June 12, Pallas, Georgia, R. Venable (**4 positives! Widest separation**)
- 2007 Jan. 10, Nysa, Georgia, R. Venable
- 2007 Feb. 21, Thisbe, Florida, D. Dunham
- 2007 Feb. 28, Nemausa, California, D. Dunham
- 2007 Apr. 13, Fortuna, Virginia and N. Carolina, D. Dunham (2 +, 1 miss, my widest separation)
- 2007 Apr. 22, Dike, Florida, R. Venable
- 2007 May 24, Papagena, Maryland and Pennsylvania, D. Dunham (3 positives)
- 2007 Sept. 11, Senta, New South Wales (AU), D. Gault (**first outside USA**)
- 2007 Nov. 20, Amalia, Georgia, R. Venable
- 2007 Dec. 18, Thusnelda, Florida, D. Dunham
- 2008 Jan. 14, Sicilia, Alabama, R. Venable (star close double)
- 2008 Feb. 10, Dynamene, North Carolina, R. Venable
- Many other cases where 2 stations were run and 1 had an occ’n & the other a miss, especially by Roger Venable; example was my observation of Rhodope occulting Regulus on 2005 October 19

(99) Dike 2005 Dec 1 $80.6 \pm 0.8 \times 61.5 \pm 0.7$ km PA 87.2 ± 1.7
 Geocentric X 4372.2 ± 0.4 Y 1805.0 ± 0.3 km N
 Double : Sep $1.1 \pm 0.3^*$, PA $30.3 \pm 19.7^*$



Center X	-1.4	0.0	✓
Center Y	-7.0	0.0	✓
Major Axis (km)	80.6	0.0	✓
Minor Axis (km)	61.5	0.0	✓
Orientation	87.2	0.0	✓
Seprn (masec)	1.1	0.0	✓
PA of 2nd star	30.3	-0.1	✓
Include Miss events		✓	
Circular	<input type="checkbox"/>	Quality	Good
Plot Scale		rms Fit: 0.2 ± 1.4 km	
<input checked="" type="radio"/> Plot both stars <input type="radio"/> Primary star only <input type="radio"/> Secondary star only			
1	David & Joan Dunham, Greenbelt, MD		
2	David & Joan Dunham, Greenbelt, MD		
3	Joe Sedlack, La Plata, MD		
4	David Dunham (mobile), Dahlgren, VA		
5 (M)	Wayne Warren, Goddard Optical Site, MD		
6 (M)	Gary Fishkorn, Dayton, MD		
7	John Wetmore, Bethesda, MD		
8	David Dunham (remote), La Plata, MD		
9	David Dunham (remote), La Plata, MD		
10	David Dunham (mobile), Dahlgren, VA		

**Multi-Station Occultation
Observing with Galileo Sized
Optical Systems**

Scott Degenhardt, IOTA

**Galileo's Legacy 2009
Waianae, Hawaii**



The Mighty Mini

Introduced to IOTA Aug 21st, 2008



50mm objective, f/2 effective f/ratio (with Owl focal reducer)


Complete portable occultation timing setup (air carryon)



Taken before boarding my plane for (343) Ostara, 7 complete stations as a carryon!



Hertha model from light curve data



Still the record, Scotty observed from 14 stations in Oklahoma

(135) Hertha 2008 Dec 11 $96.5 \pm 2.3 \times 64.3 \pm 2.5$ km, PA 40.6 ± 3.0
Geocentric X 3159.2 ± 0.9 Y 1940.4 ± 1.1 km

N

E

Find best fit

☒ Center X ☒ -6.7
☒ Center Y ☒ 3.4

Major axis (km) ☒ 4.3
Minor axis (km) ☒ -6.7
Orientation ☒ 5.0

a/b=1.50
dM=-0.44

Double star
Sepn (masec) ☐ 0.0
PA of 2nd ☐ 0.0

☒ Both ☐ Primary ☐ Secondary


☐ Circular ☐ Include Miss events

Plot scale Quality

RMS fit 1.8 ± 7.5 km

1 (M)	R Stanton, Three Rivers, CA
2 (M)	P Maley, Baker CA
3	G Mroz, Santa Fe, NM
4	S Degenhardt01, Okarche, OK
6	D Dunham, Harvard, CA
7	S Degenhardt02, Okarche, OK
8	S Degenhardt03, Okarche, OK
9	S Degenhardt04, Okarche, OK
10	R Wasson, Barstow, CA
11	S Degenhardt05, El Reno, OK
12	S Degenhardt06, El Reno, OK
13	S Degenhardt07, El Reno, OK
14	S Degenhardt08, El Reno, OK
15	S Degenhardt09, El Reno, OK
16	A Holmes, Goleta, CA
17	S Degenhardt10, El Reno, OK
18	S Degenhardt11, El Reno, OK
19	S Degenhardt12, Union City, OK
20	S Degenhardt15, Union City, OK
21	S Degenhardt14, Minco, OK
22	B Owen/J Young, Wrightwood, CA
23	K Young, Wrightwood, CA
24	G Lyzenga, Altadena, CA

Number of extra observing stations above 1 per person 2000-2009



Mighty Mini



Can record
occultations of stars
to mag. 9.5, even
mag. 10.0 under
good conditions

Mighty Midi – Orion 80mm short tube



Can record
occultations of stars
to mag. 11.0, even
mag. 11.3 under
good conditions

I use visual finder
scope and \$60
Quantanray tripod
while Scotty uses a
mighty mini video
as the finder and
MX-350 tripod
(not as sturdy as the
Quantanray)

Mighty Maxi – Orion 120mm short tube




Can record occultations of stars to mag. 12.0, even mag. 12.5 under good conditions


I use visual finder scope and an alt-az mount built from PVC pipe, bolts, wing nuts, velcro straps, 2 “clam shells” made from large-diameter PVC pipe, and a rectangular piece of wood that Scotty sold me for \$50. The 2-pt. support for the heavy scope make balance and altitude adjustment tricky; placing an MX-350 tripod under the end of the camera with crumpled paper between adds a 3rd point and stability, but re-pointing is often needed when put in place. Scotty has a better mount design for about \$100 in parts that he will present at the IOTA meeting in Oct. Commercial mounts that can hold this weight cost hundreds of \$, more than twice the \$300 cost of the 120mm OTA.

Scotty's Maxi Mount

- Solid as a rock
- All sky accessible
- Air portable @ < 12 lbs.
- (<20 including scope)
- 2 axis slow motion control
- Stealthy black for multi-station deployment
- Stands only 24" tall
- Costs ~\$100



Programmable Remote for Timed Recordings



Suggested by Steve Conard. Scotty found a “100% effective” system. Place transparent plastic tube (I believe made from 2 coin holders fastened together with Scotch tape; shown at foot of tripod) at bottom of the brown mailing bag in the background. After setting the programmable remote, place it pointing down at the tube at the bottom. Turn the Canon ZR camcorder to the VCR position with front end down facing the tube. If cold, add some hand warmers. 6 plastic tabs glued to the edges of the front of the remote, and the piece of cardboard held on with the rubber band, prevent the programmable remote from turning on, which happens whenever the screen is touched.

Components of
John Broughton's
25cm "Suitcase
Telescope"



Goal:
Fit 2 of them
within the
22-kg per
suitcase
weight limit
of most
airlines




The Suitcase Telescope set up for observing (except for the cameras).
3 can be set up ready to go and put on the back seat area of an ordinary car; more could be put in the trunk.




Occultation of 7th-mag. Star by (234) Barbara, 2009 Nov. 21




Path of the 2010 July 8th occultation of 2.5-mag. Yed Prior (delta Ophiuchi) by the asteroid (472) Roma




Sky chart for the 2010 July 8th occultation of 2.5-mag. Yed Prior (delta Ophiuchi) by the asteroid (472) Roma



Pre-Point chart for the 2010 July 8th occultation of 2.5-mag. Yed Prior (delta Ophiuchi) by the asteroid (472) Roma




Path of the 2010 July 8th occultation of 2.5-mag. Yed Prior (delta Ophiuchi) by the asteroid (472) Roma over Iberia




Mighty Mini Training at Sabadell



Path of the 2010 July 8th occultation of 2.5-mag. Yed Prior (delta Ophiuchi) by the asteroid (472) Roma over s.w. Iberia




Locations of Mighty Minis deployed for the 2010 July 8th occultation of Yed Prior by Roma in s.w. Iberia




Our First Station, Setup in a ditch at La Albuera, Spain, s.e. of Badajoz




Saturation study of Portuguese mighty mini light curves



Saturation study of Gonçalves' mighty mini light curve



Joao Cruz' unsaturated partial occultation light curve



695 Bella occults TYC 2332-01054-1 on 2010 Aug 31 from 12h 5m to 12h 29m UT

Star:

Mv = 7.7 Mp = 7.8 Mr = 7.6

RA = 2 33 42.7562 (J2000)

Dec = 34 20 28.608

[of Date: 2 34 24, 34 23 19]

Prediction of 2010 Jul 21.0

Max Duration = 5.4 secs

Mag Drop = 5.5 (5.1x)

Sun : Dist = 110 deg

Moon: Dist = 16 deg

: illum = 62 %

E 0.028"x 0.019" in PA 87

Asteroid:


Mag = 13.2

Dia = 48km, 0.041"

Parallax = 5.500"

Hourly dRA = 1.504s

dDec = 20.23"







Occult Watcher Stations for the Bella Occultation




Successful Stations for the Bella Occultation



Bella Station 9 reappearance shows that the star is likely a close binary




Bella Station 10, shows a step reappearance like Station 9



Profile of Bella from the Aug. 31st Observations

Stations mostly set up s. of predicted center due to expected PPMXL star catalog shift



Bella Occultation my best success, but . .

- It was extremely exhausting, made me ill
- As I drove back to Fresno, my nose started running
- At hotel, 2 hours to pack, then go to airport – no rest
- Ear infection evident as plane landed in Baltimore
- Next week, very weak, flu symptoms for 3 days
- Didn't recover; after 2 weeks, I saw doctor
- Diagnosis: Thyroiditis; main recovery took 6 weeks
- Full recovery took several months
- Now, I insist on 4h+ sleep after an all-nighter
- Need younger people to take up this extreme sport!

Occultation of LQ Aquarii on
2011 July 19
mapped the profile
of the binary asteroid (90) Antiope

David W. Dunham,
International Occultation Timing Association
and KinetX, and many others, from IOTA,
IMCCE (Paris Observatory), the SETI Institute,
Southwest Research Institute, JPL, Sierra College,
etc.


90 Antiope occults HIP 112420 on 2011 Jul 19 from 10h 13m to 11h 6m UT

Star: Dia = 2mas
Mv = 6.7 Mp = 8.3 Mr = 5.8
RA = 22 46 14.2117 (J2000)
Dec = -11 9 59.068
[of Date: 22 46 53, -11 6 6]
Prediction of 2011 Jun 8.0

Max Duration = 40.4 secs
Mag Drop = 5.8 (6.2r)
Sun : Dist = 138 deg
Moon: Dist = 10 deg
illum = 83 %
E 0.030"x 0.024" in PA 78


Asteroid:
Mag = 12.5
Dia = 120km, 0.091"
Parallax = 4.811"
Hourly dRA = -0.421s
dDec = -5.47"

Expect fades - star dia. Variable star Asteroid has 1 moon(s). 84km at 171km



Predicted Path of the
Occultation of the
6.7-mag. Star by
(90) Antiope

The path over northern Calif. & Nevada





(90) Antiope 2011 Jul 15 15:08:13 UT

Keck AO Kp-band, exp 3 sec

rotation sense is CCW

31 min plus 5.5 cycles prior to occultation

Merline, Neyman, Tamblyn, et al.



PA 189 deg, projected separation 146 km



0

2000

4000

6000

8000

10000

12000


14000


16000

IOTA meeting at Sierra College, Rocklin, CA



One of the pre-point charts prepared for
the 50mm binocular lens-based
“mighty mini” systems






Setting up a mighty mini at my station #5 in Newman, Calif.




Light curve of the occultation at Dunham Station #1 west of Tracy, Calif.




The angular diameter of the red giant star caused the disappearance and reappearance to be gradual over several tenths of a second, with different durations at the two events due to different slopes of the asteroid's surface

Preliminary Sky-plane Profile




I ran 10 stations in Florida for the occultation of 4 Tauri by Hertzprung July 30 am



My stations crossed both
1-sigma zones from
40 mi. s.e. of Orlando
20 mi. w. of W. Palm
2 failed (not shown)
2 recorded the occultation
(green circles)
6 had no occultation
(red circles)
Three others observed
from home north of
my observing fence
Actual path was in n.
1-sigma zone, as
predicted by PPMX
Spectral class A star v

Hertzsprung Event Sky-plane Profile




407 Arachne occults HIP 54719 on 2011 Dec 28 from 11h 17m to 11h 41m UT

Star:
 Mv = 8.2 Mp = 8.4 Mr = 8.1
 RA = 11 12 10.3714 (J2000)
 Dec = - 0 24 34.660
 [of Date: 11 12 49, - 0 28 39]
 Prediction of 2011 Dec 15.0

Max Duration = 11.9 secs
 Mag Drop = 5.7 (5.4r)
 Sun : Dist = 107 deg
 Moon: Dist = 154 deg
 : illum = 16 %
 E 0.030"x 0.019" in PA 98


Asteroid:
 Mag = 13.9
 Dia = 98km, 0.059"
 Parallax = 3.837"
 Hourly dRA = 0.730s
 dDec = -13.90"




Arachne OW stations 3-9



Arachne OW stations 1 & 2 (cloudy)



Arachne Sta. 3, Foster Cemetery




locations currently announced by other observers:

US287 e. remote; 2 = Iverson ; 3 = Lowe A Home; 4 = Glendale TX remote; 5 = Trinity TX remote; 6 = Davis Hall Rd remote; 7 = e Huntsville TX remote; 8 = w Hur


ntly selected site coordinates are: $-95^{\circ} 45' 51''$, $30^{\circ} 40' 51''$. The site is 43 km left (NE) from the central line.

Sky-Plane Plot



Limovie Lightcurve for my Station #3 at Foster Cemetery, e. of Huntsville, TX

Analyzed file name [20111228_Arachne_Dunham3b V1.AVS] Photometry in each Frame



Analyzed by Scotty
Dunham

Occultation of SAO 60804, mag.
8.0, by the Trojan Asteroid
(911) Agamemnon
Thurs. morning, 2012 January 19

This is the best, most valuable asteroidal occultation over
the MD/greater Washington, DC region this year

We need your help to observe it

Following are maps of the path, and several star charts to
locate the star in obscure Lynx;

**An article about the results of this occultation is in
press for publication in Planetary and Space Science**

911 Agamemnon occults HIP 41337 on 2012 Jan 19 from 11h 31m to 11h 41m UT

Star:
Mv = 8.0 Mp = 9.0 Mr = 7.5
RA = 8 26 2.3658 (J2000)
Dec = 36 58 57.397
[of Date: 8 26 52, 36 56 21]
Prediction of 2011 Dec 16.0

= SAO 138052,


spec. type K0

Max Duration = 10.4 secs
Mag Drop = 6.8 (6.9r)
Sun : Dist = 163 deg
Moon: Dist = 128 deg
illum = 17 %
E 0.046"x 0.024" in PA 115

Occultation in Mid-Atlantic
region at 11:31:40 UT $\pm 11s$
(6:31am & 40s EST)


Asteroid:
Mag = 14.8
Dia = 185km, 0.062"
Parallax = 2.137"
Hourly dRA = -1.789s
dDec = 0.83"

In coastal Mid-Atlantic
region (DC, Philadel-
phia, Richmond, etc.),
Star alt. 24° in az. 299°
(northwest, opposite
Sun, whose alt.
will be -10°)



Finder chart, 2° on a side


Map of 2012 Jan. 19th Agamemnon occ'n observers




Dunham ran a 2nd station between #1 and #3 but the video recorder at #2 failed.

Green dots mark positive observations, while red (Brooks) indicates a miss (negative, no occ'n). The green line is the predicted central line while the blue lines are the predicted path limits. The gray lines mark a 10-km zone bracketing Conard's location where the satellite occ'n occurred.

Steve Conard's light curve of the Jan. 19th Agamemnon occultation



Sky-plane plot, 2012 Jan. 19th (911) Agamemnon occultation




Agamemnon: Axes 190.6 ± 0.9 by 143.8 ± 1.5 km, PA minor axis $-69.3^\circ \pm 1.3^\circ$,
center X 4661.6 ± 0.4 km, Y 3113.7 ± 0.6 km; disappearances on right side
Satellite plotted as 9-km circle (but it's more likely about 4 km across)
 $0.0931''$ (278 km in the plane) from Agamemnon's center in PA 93.8°

Occultation of 6.5-mag. SAO 140947 (HIP 78870) by asteroid (28) Bellona in the Khabarovsk area, 11 May 2012 evening

A plan to observe the occultation with five “mighty mini”s
set up across the path to measure the size and shape of Bellona

**David W. Dunham, KinetX, Inc.,
Moscow Inst. of Electronics & Mathematics (MIEM),
and
International Occultation Timing Association (IOTA)**

The best previously-observed Bellona occultation was observed in the northeastern U.S.A. almost 10 years before




Next good Russian asteroidal occultation, Khabarovsk area, 11 May 2012

28 Bellona occults HIP 78870 on 2012 May 11 from 12h 33m to 12h 46m UT


Star:
Mv = 6.5 Mp = 7.4 Mr = 6.0
RA = 16 5 59.7900 (J2000)
Dec = - 6 8 23.207
[of Date: 16 6 42, - 6 10 25]
Prediction of 2012 Apr 19.0

Max Duration = 10.7 secs
Mag Drop = 4.7 (4.7r)
Sun : Dist = 163 deg
Moon: Dist = 63 deg
illum = 65 %
E 0.026"x 0.022" in PA 95


Asteroid:
Mag = 11.2
Dia = 128km, 0.094"
Parallax = 4.705"
Hourly dRA = -2.065s
dDec = 8.05"




Naked Eye View, towards the southeast




80mm Short-Tube Refractor or “Mighty Midi” view, (28) Bellona
and SAO 140947, Khabarovsk area, 11 May 2012



80 коротких труб рефрактор или â € € œMighty Мидия зрения (28),
Беллона и CAO 140947, Хабаровский район, 11 мая 2012



Прямоугольник показывает могучий миди области
просмотра с целью звездой в центре









Mighty Mini Training in Khabarovsk






Stations around Khabarovsk for the 2011 May 11 (28) Bellona Occultation



Observing locations currently announced by other observers:


3 = Fleishman R Home; 4 = Vladimirovka Igor dacha; 5 = 7km e. Sita, RU; 6 = Sukpay, RU ; 7 = David Dunham (100km nne KHV friend dacha)

Occultation of 9.5-mag. SAO 185777 by (2258) Viipuri observed by RECON Team



The RECON Team is a 12-station initial system of 11-in. SCT's set up by the South-West Research Inst. at colleges and high schools in small towns in the good-weather rain shadow of the Cascade and Sierra Nevada mountains; the full system may extend across the USA from the borders with Canada and Mexico. for TNO's, but practice with MBA Events.

Satellite of (2258) Viipuri apparently found




Occultation of 10.6-mag. TYC 0030-00604-1 by (307) Nike, 2013 October 6, during IOTA meeting

307 Nike occults TYC 0030-00604-1 on 2013 Oct 6 from 2h 44m to 2h 55m UT

Star:
Mv = 10.6 Mp = 11.4 Mr = 10.2
RA = 1 52 41.5260 (J2000)
Dec = 1 12 41.285
[of Date: 1 53 26, 1 16 52]
Prediction of 2013 Aug 13.0

Max Duration = 6.2 secs
Mag Drop = 3.2 (3.1r)
Sun : Dist = 163 deg
Moon : Dist = 169 deg
illum = 1 %
E 0.044"x 0.027" in RA 73

Asteroid:
Mag = 13.7
Dia = 60km, 0.051"
Parallax = 5.439"
Hourly dRA = -1.793s
dDec = -12.04"



Occultation of 9.6-mag. TYC 0646-00730-1 by binary Trojan (617) Patroclus, 2013 October 21


617 Patroclus occults TYC 0646-00730-1 on 2013 Oct 21 from 6h 38m to 6h 50m UT

Star:
Mv = 9.6 Mp = 11.3 Mr = 8.7
RA = 2 47 58.7769 (J2000)
Dec = 14 5 59.732
[of Date: 2 48 46, 14 9 28]
Prediction of 2013 May 8.0

Max Duration = 8.6 secs
Mag Drop = 5.1 (5.6r)
Sun : Dist = 164 deg
Moon: Dist = 11 deg
illum = 95 %
E 0.084"x 0.033" in PA 56

Asteroid:
Mag = 14.7
Dia = 141km, 0.052"
Parallax = 2.370"
Hourly dRA = -1.614s
dDec = 0.42"

Variable star Asteroid has 1 moon(s), 113km at 680km



Occult4103

Occultation of 6.7-mag. SAO 162945 by (41) Daphne, 2013 October 25

41 Daphne occults HIP 97157 on 2013 Oct 25 from 12h 30m to 12h 34m UT

Star: Dia = 2mas
Mv = 6.7 Mp = 8.3 Mr = 5.8
RA = 19 44 54.2992 (J2000)
Dec = -10 34 23.233
[of Date: 19 45 40, -10 32 9]
Prediction of 2013 Jul 8.0

Max Duration = 9.0 secs
Mag Drop = 5.6 (6.0r)
Sun : Dist = 84 deg
Moon: Dist = 169 deg
: illum = 64 %
E 0.015"x 0.009" in PA 89

Asteroid: (in ISAM)
Mag = 12.3
Dia = 210km, 0.116"
Parallax = 3.515"
Hourly dRA = 3.133s
dDec = -7.69"

Expect fades - star dia. Asteroid has 1 moon(s), 2km at 443km




Occultation of Regulus by (163) Erigone, 2014 March 20

163 Erigone occults HIP 49669 on 2014 Mar 20 from 5h 53m to 6h 22m UT

Star: Dia = 1mas
Mv = 1.3 Mp = 1.3 Mr = 1.3
RA = 10 8 22.0688 (J2000)
Dec = 11 58 2.038
[of Date: 10 9 9, 11 53 37]
Prediction of 2013 Apr 10.0

Max Duration = 14.3 secs
Mag Drop = 11.1 (10.7r)
Sun : Dist = 150 deg
Moon: Dist = 72 deg
illum = 87 %
E 0.029"x 0.013" in PA 105

Asteroid:
Mag = 12.4
Dia = 72km, 0.084"
Parallax = 7.421"
Hourly dRA = -1.110s
dDec = 13.72"



Since the Occultation by (166) Rhodope in Oct. 2005, Regulus was found to be a close double, but the companion is about 100 times fainter than the primary, and it apparently was not detected in the 2005 observations. I will show my remote Video.